



## King's Research Portal

DOI:

[10.1016/j.jceh.2019.05.005](https://doi.org/10.1016/j.jceh.2019.05.005)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Fitzpatrick, E., & Dhawan, A. (2019). Childhood and Adolescent Nonalcoholic Fatty Liver Disease: Is It Different From Adults? *Journal of Clinical and Experimental Hepatology*, 9(6), 716-722.

<https://doi.org/10.1016/j.jceh.2019.05.005>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## **Childhood and adolescent NAFLD: IS it different from adults?**

Emer Fitzpatrick and Anil Dhawan

Paediatric Liver, GI and Nutrition Centre, King's College London Faculty of Life Sciences and  
Medicine and King's College Hospital

### ***Corresponding author:***

Professor Anil Dhawan

Paediatric Liver, GI and Nutrition Centre

King's College Hospital

Denmark Hill

London SE5 9PJ

Tel: +442032994408

[anil.dhawan@kcl.ac.uk](mailto:anil.dhawan@kcl.ac.uk)

Neither author has any conflict of interest to declare

## **Abstract**

Paediatric non-alcoholic disease (NAFLD) is the most common chronic liver disease in childhood and adolescence. Though the condition is similar in many ways to NAFLD in adults, there are important differences in predisposition, presentation, differential diagnosis and potentially also in optimal management. Antenatal and early childhood exposures and the particular vulnerabilities in a growing child to environmental influences present unique opportunities for intervention and modification of risk. The prevalence of significant fibrosis on biopsy in pre-adolescent children in the context of NAFLD should not be ignored but the relevance of this fibrosis to long term outcome is as yet unknown. Approach to children and adolescents with suspected NAFLD needs to include an assessment of risk factors in addition to exclusion of alternative or co-existing liver diseases. Liver biopsy is indicated in younger children and in those without clear predisposing factors leading to metabolic syndrome, also in those in whom significant fibrosis is suspected. The histology in children and adolescents differs from adults in that type 2 NAFLD may be more prevalent, which is associated in turn with more significant fibrosis. Management in children and adolescents needs to focus on lifestyle intervention, which, when weight loss is achieved, demonstrates excellent results in terms of resolution of disease. Appropriate intervention in childhood and adolescence where may prove instrumental in avoiding the need for later transplantation while also decreasing all-cause mortality in these at-risk individuals.

**Key words:** Paediatric NAFLD, early-life course, risk factors, management

## **Introduction**

NAFLD is rapidly becoming the most common indication for liver transplantation liver in the western world (1). Paediatric NAFLD is thought of as an early presentation of the adult condition and in general is thought not to frequently progress to decompensated end stage liver disease nor hepatocellular carcinoma in childhood. That said, paediatric NAFLD presents both challenges and opportunities in terms of both diagnosis and management. Several important distinctions exist between adult and paediatric disease which need to be understood before we can effectively approach management. It is not clear why some patients with NAFLD present during childhood with often significant fibrosis; it is possible that this is merely incidental and the course of disease is still over 40 – 50 years prior to end stage. The alternatives, however, are that paediatric NAFLD is a distinct disease with different susceptibilities and pathophysiology than the adult disease or that presenting in early life forebodes more significant or severe disease (2). The question of the long-term outcome of those with paediatric-onset NAFLD is as yet unanswered.

In the first instance, early life susceptibilities to the condition are most often identifiable in paediatric versus adult onset NAFLD. Preconception maternal obesity and gestational diabetes are recognised as important risk factors which need to be addressed at a societal and preventative level (3). Early feeding practices, weaning and the exposure of children to the dangers of the high sugar westernised diet particularly during a vulnerable stage of development are all relevant. The rapid physical and psychological changes that occur during adolescence where insulin resistance peaks and body mass is laid down are critical periods where metabolism is regulated and may also be important in establishing or avoiding liver damage.

This review will approach paediatric NAFLD from the viewpoint of a comparison to adult disease and the different approach that may be taken in light of this distinction.

## **Epidemiology**

NAFLD is thought to affect 10% of children under the age of 18 years as demonstrated by a post mortem study of livers of children and young people who suffered unnatural death. Steatosis was found in 9% of children with 3% having evidence of liver inflammation / fibrosis (4). Epidemiological studies are less likely to use histological definitions of the disease and the prevalence varies accordingly. When ultrasound is used to assess steatosis, prevalence varies from 1.8% in a normal population of children and young people to 60% of children and young people undergoing bariatric surgery to 80% of those in an obesity clinic (5-7).

These figures are not far from adult prevalence and are closely linked to the prevalence of obesity. Severity of disease in children does not necessarily associate directly with severity of obesity for example and possibly different prenatal and childhood exposures on a genetically susceptible individual may predispose to early onset of disease. The degree of obesity in an individual does not correlate to severity of liver disease in that individual(8). The concept that those with certain genetic susceptibilities should have a 'normal' BMI which may be less than the conventional 'normal' holds true for this condition.

### **Susceptibility**

The effects of the intrauterine environment in terms of priming may be more relevant to paediatric patients in terms of developing NAFLD (9). Both small for gestational age and large for gestational age infants are overrepresented in those who develop NAFLD during childhood and adolescence(10). Intrahepatic lipid content investigated using MRI is higher in infants of mothers who were obese and those with type 2 diabetes than in those born to normal weight mothers (11, 12). Interestingly intrahepatic fat content was more closely related to maternal BMI than to birthweight (12). In a post mortem study of stillborn infants, those born to mothers with gestational diabetes had a prevalence of 78.8% steatosis versus those still born infants of mothers who were not diabetic (13). Deposition of subcutaneous fat does not occur until the third trimester this it is conceivable that there is hepatic storage of excess substrate in the fetal liver, in addition to in utero de novo lipogenesis in response to a high transplacental glucose supply (9). Insulin does not cross the placenta but it is thought to increase placental inflammation altering the transfer of nutrients to the fetus. In addition, increased fatty acids signal placental Toll like receptor 4 expression, again this proinflammatory response may increase fetal nutrient transport (14). Placental insufficiency has also been associated with a higher inflammatory milieu and increased metabolic risk (15). There is a clear relationship between maternal obesity and BMI in childhood (16, 17). It is thus likely that the consequent complications of obesity including NAFLD will follow this increased risk (18).

The link between antenatal and early postnatal exposure and NAFLD is complex and multifactorial. In part the possibility that the microbiome may be involved in establishing susceptibility to poor metabolic health. Infants with decreased microbiome diversity at 6 months are greater risk obesity aged 7 years (19). It is known that immune tolerance is promoted by gut microbiota, this is also decreased in number in offspring of obese women . Breast feeding, which has shown to be protective against NAFLD in some studies (though the confounding principles of socioeconomic influences on this observation is difficult to unpick) may act in part at least via the microbiome, though the confounding issue of socioeconomic influence . Breast feeding promotes colonisation of the intestinal microbiome providing oligosaccharides as prebiotics(18).

Animal studies can elucidate further the antenatal risk to the fetus and the compounding influence of early life exposure. In mice, a study reported the influence of high fat diet (HFD) in dams and in pups versus controls. HFD in dams led to fatty liver in offspring, this was seen to a greater extent when the pups were also fed HFD but also seen in normal fed chow fed pups. Both high fat diet pre and post pregnancy leads to a cumulative risk (20). Both DNA methylation alteration (21) and a decrease of diversity in the microbiome are both possible mediators of this effect. Another mouse model of maternal obesity demonstrated higher oxidative stress and impairment of innate immunity in pups following an in utero high fat diet. At 12 months the pups showed steatohepatitis and fibrosis with an increased number of inflammatory and fibrogenic mediators (22).

A similar experiment in rats demonstrated a sex prediction in that male offspring of high fat dams demonstrated more injury than females; possibly due to the different growth trajectory expected in male versus female pups (23).

Non-human primates fed a high fat diet prior to breeding which was then normalised during pregnancy demonstrated that effects on offspring can be modified. An increased liver triglyceride content with increased expression lipogenic genes in liver tissue and increased activation of inflammatory gene expression was demonstrated in these animals (24).

## **Genetics**

In terms of genetic susceptibility, findings in children largely mirror those in adults with the minor allele of PNPLA3 (rs738409) widely reported as a susceptibility factor (25, 26) and frequency varies according to ethnicity of the patients affected. Other genetic variants such as TM6SF2 rs58542926 (27) (28) and GCKR (29) are implicated in susceptibility but there have been few widespread paediatric GWAS studies and only one in biopsy proven- NAFLD in children(30). In this study 234 Hispanic boys were investigated for genetic variations predisposing them to NAFLD. In addition to PNPA3 and TMSF6, the authors found that novel variants in Trafficking Protein Particle Complex 9 (TRAPPC9) were associated with the NAFLD activity score (NAS) and a single nucleotide polymorphism (SNP) in a region close to ARP5 – actin related protein 5 was associated with fibrosis.

Clearly genetic variation plays some part but is not in itself sufficient to explain propensity to disease. Given that even within the obese population, genetic susceptibility only comprises part of the risk, much focus has been given to dietary components and whether specific diets convey risk.

### **Dietary intake**

Many studies have drawn associations between the intake of fructose, saturated fat and decreased fibre intake, polyunsaturated fats and the development of NAFLD (31). Children have a particularly high consumption of fructose containing sweetened beverages up to 300kcal / day (32). In addition children have the highest intake of ultra-processed food at 33% (32.1-35%) versus adults 29.6% (28.5 – 30.7%) (33). Both have been linked to poor metabolic health including NAFLD. Indeed, the consumption of a Western diet was strongly associated with NAFLD in both Australian (34) and Chinese adolescents (35). A review of children with biopsy proven NAFLD compared to obese controls in the UK did not reveal any major dietary differences between the two groups however the NAFLD group tended to be lighter and more active in general (36). It is difficult to conclude that inclusion or exclusion of one or more component of diet may make a difference to the development and progression of NAFLD in children.

### **Diagnosis**

The diagnosis of NAFLD in children brings challenges. Though significant alcohol consumption particularly in children less than 14 years is unlikely to play a major contribution to liver injury many other liver diseases of

childhood may present with steatosis with or without inflammation and fibrosis. Given that 30% of the paediatric population worldwide is overweight or obese (37), in those presenting with Wilson disease or other Hepatitis C the presence of overweight and obesity equates to the population norm. Thus, the presence of overweight or obesity does not mean that a child with steatosis does not have an alternative diagnosis to NAFLD! The work up for fatty liver, usually found incidentally when a child has blood tests for another reason and goes on to have an ultrasound scan due to abnormal LFTs, should be comprehensive (38). In addition to screening for associated features of the metabolic syndrome such as fasting lipid levels, hypertension, HbA1C, HOMA-IR and impaired glucose tolerance, children should undergo a work up for alternate liver conditions. This should include but is not limited to infectious hepatitis, Wilson disease, inborn errors of metabolism (fatty acid oxidation disorders, mitochondrial disease), coeliac disease, alpha 1 antitrypsin deficiency and abnormalities of lipid metabolism (for example; hypobetalipoproteinemia). There are several other conditions that may be investigated based on clinical suspicion and summarised in Table 1. In addition, a full family, feeding and medication history should be taken. Parenteral nutrition and medications such as steroids, antipsychotics and antidepressants may all predispose to weight gain and steatosis.

Though the definition of NAFLD is a histological one, non-invasive methods to detect and stage the disease are now common in clinical practice (39). The most useful is ultrasound which can detect > 30% steatosis which is probably the level at which it is most clinically significant. ALT, AST and GGT levels may reveal liver inflammation at a point in time but will not reliably differentiate those with and without fibrotic disease which is the main determinant of outcome.

Radiological techniques such as MRS and MRI proton dense fat fraction will differentiate different grades of intrahepatic lipid and MR elastography is a useful measure of liver stiffness – a proxy for fibrosis (40, 41). MRI is expensive and cumbersome however and not suited to day-to-day clinical practice though can be a useful research tool.

Transient elastography is well validated in paediatric as well as adult studies to detect fibrosis in NAFLD (41, 42). The relatively new controlled attenuation parameter measurement may also prove useful in quantifying steatosis in a longitudinal manner but use in children has not been adequately validated as yet (43).

Acoustic radiation force imaging and other types of shear wave elastography (aside from TE), are other emerging methods of detection and quantification of fibrosis (44). They have not yet been extensively validated in paediatric patients with NAFLD though encouraging reports are emerging (45).



Such is the enormity of the prevalence of NAFLD in the population, the question of who and when to refer to the paediatric hepatologist is a difficult one. In general, in those with a fatty liver on ultrasound, NAFLD is a diagnosis of exclusion. In the absence of the typical phenotype, other conditions must be suspected first. Even in those with other features of the metabolic syndrome and a relevant family history of NAFLD, other or co-existing disease should be suspected. Severity of disease is not easy to elucidate in primary care. Several algorithms have been developed in large adult cohorts to differentiate significantly fibrotic disease. Unfortunately, most are not applicable to children given that they include age, BMI (adult reference) or markers of collagen turn over. For example, it is well established that FIB4 is unreliable in those under the age of 35 years (46), the fatty liver index uses BMI, the absolute number of which is not applicable to children (rather need BMI z score or centile) (47). ELF score uses P3NP as a variable which is an age and sex dependent marker of collagen turn over.

The Pediatric NAFLD score is the only paediatric-specific algorithm, but still has not been validated outside of a Caucasian Italian population (48).

Thus, as per both ESPGHAN and NASPGHAN guidelines, we first need to be cautious about making a positive diagnosis of NAFLD, considering relevant differential diagnosis (49, 50).

Histological diagnosis is ideal though only rarely practical. Assessing severity of disease is focused on detection and staging of fibrosis. Until such time that appropriate serum biomarkers or algorithms are available and validated, the non-invasive determination of degree of fibrosis is still limited to larger centres which have transient elastography or other imaging techniques for fibrosis readily available.

## **Histology**

Typical 'paediatric' NAFLD differs histologically from that in adults (51). This pattern, found predominantly in children and young people has been labelled 'type 2 NAFLD' and is periportal in distribution versus the largely pericentral disease in adults. In various series the prevalence of type 2 histology differs (51, 52). Often, children may have a mixed type 1 and type 2 pattern. In type 1 NAFLD, steatosis, inflammation and fibrosis are mainly lobular, surrounding the central vein. In type 2, steatosis, inflammation and fibrosis are preferentially periportal. Ballooning, one of the classical histological features of NAFLD, is less common in children with type 2 disease

Table 3 – differences

The NAFLD activity score (NAS) developed by a collaboration of pathologists using material from adults and children with NAFLD through the NASH clinical research network (53). It is the most commonly used scoring system in the literature and is based on a score of 0 – 3 for steatosis, 0 – 2 for ballooning and 0 – 3 for lobular inflammation. Fibrosis is scored separately from 0 to 4. The diagnosis of non-alcoholic steatohepatitis from a research perspective is based on a NAS score of 5 or more. A score of 2 or less is not NASH and 3 or 4 is borderline. Children tend to score lower on the NAS due to less lobular inflammation and ballooning. The presence of NASH was previously thought to be the most important prognostic factor in determining outcome in patients with NAFLD. It is now recognised however that the presence of fibrosis is a more reliable predictor of outcome, not least as the degree of inflammation in a biopsy may change over days to weeks (54).

The presence of type 2 disease is associated with more advanced fibrosis and progressive disease (55). Steatosis may be macro or microvesicular or both. The presence of microvesicular steatosis should prompt the possibility of mitochondrial disorders or other inborn errors of metabolism, though the prevalence of microvascular steatosis in adult onset NAFLD is reported and associated with greater severity of disease.

The reason that children demonstrate relatively greater prevalence of type 2 disease is not entirely clear. We speculate however that it may possibly be linked to the development of zonation in the liver lobule (56, 57). Enzymes involved in Krebs's cycle for example found mostly in zone 1 in the periportal region. Hepatocytes responsible for detoxification and metabolism due to the p450 mechanism are preferentially located in zone 3 (perivenular). It is possible that the differential function along the lobule may predispose children to susceptibility to injury at different stages of development (56, 57).

### **Natural history and treatment**

A handful of case series have reported on the long-term follow up of children with NAFLD including the need for liver transplantation at an early age, though be it rare (58-60). The difficulty with assessment of severe paediatric fatty liver disease is the concern that an alternative undiagnosed metabolic disease may be present, particularly in the case of 'lean NAFLD' (38). In adults it is thought to take approximately 7 years to progress one fibrosis stage (60). A study analysed paired liver biopsies from 122 children who were in the placebo group in two randomised clinical trials for NAFLD (lifestyle advice only for either 52 or 96 weeks). During the trial period,

the fibrosis stage progressed in 23%, and improved in 34%. Younger children with more severe fibrosis at time of biopsy may progress more readily and it is likely that in these selected children, the effects of genetic variants on susceptibility may outweigh environmental factors.

Lifestyle change resulting in weight loss is an effective way of reversing or stabilising disease. A small number of trials in children have demonstrated results. In an Italian study of 84 children, weight loss average of 4kg over a 12 months period resulted in an improvement in ALT and ultrasound features of steatosis (61). There was a drop out weight of 30%. Another paediatric study of intensive lifestyle intervention in North America achieved improvement in BMI z-score with a decrease of 0.1 U ( $p<0.05$ ) baseline to one year and decrease in ALT in 69% of the follow up cohort. There was a 53% drop out rate however (62).

These results are reflected in the literature regarding adults with NAFLD; a metaanalysis of studies achieving weight loss of 5% or more resulted in improvement in steatosis whereas  $\geq 7\%$  weight loss resulted in improvement in steatohepatitis and in those with  $\geq 10\%$  weight loss, all features of NAFLD were reversed or stabilised (63). In a prospective study in adults these outcomes were confirmed (64). There was a high drop-out rate however and only 50% successfully achieved 7% weight loss or more. Of note in 94% of those who achieved  $\geq 5\%$  weight loss, fibrosis stabilised or reversed.

The success of weight loss alone on the outcome of patients with NAFLD is remarkable, yet the barriers to achieve this for all patients have not yet been adequately addressed. It is known that the co-existence of major depressive disorder, for example, is a major factor in the failure of lifestyle treatment of NAFLD. Willingness to engage in the programme and readiness to change are naturally crucial. The window of opportunity for many may be during childhood and early adolescent window as the likelihood of overweight children becoming overweight adults is  $> 80\%$ .

There is a burgeoning industry in developing drugs and compounds to treat NAFLD given the inability of many to lose weight. In children neither the TONIC trial (comparing metformin, vitamin E and placebo)(65) and the CYnCH trial (comparing cysteamine bitartrate and placebo)(66) reached their primary outcome measure. The use of vitamin E in improving some parameters of histology(ballooning) demonstrated statistical significance.

Newer, more targeted compounds such as FXR agonists, antifibrotic drugs and certain inflammation inhibitors show some promise in preclinical and early clinical trial but use in children have not yet been studied.

Bariatric surgery has been clearly shown to reverse the disease (6, 67), it is not clear whether this effect is via decreased intake / appetite control nor via other disruption of metabolic pathways by banding or by bypassing the stomach.

The opportunities to use intensive lifestyle change and maintenance of the lifestyle change through family education, counselling and an individualised approach are stark. Undoubtedly deprivation and easy availability of ultra-processed in expensive foodstuffs is impossible to tackle at an individual level and we need to exert our challenge at a societal level.

### **Conclusions**

Paediatric NAFLD is a prevalent condition world-wide. Though it shares many features with adult – onset disease, paediatric NAFLD has important differences. In particular early life influences and susceptibilities and an approach to diagnosis and management need to be considered. There is a real opportunity to reverse the course of the disease in childhood and paediatricians should be aware and ready to act.

### **References**

1. Cholaneril G, Wong RJ, Hu M, Perumpail RB, Yoo ER, Puri P, et al. Liver Transplantation for Nonalcoholic Steatohepatitis in the US: Temporal Trends and Outcomes. *Dig Dis Sci*. 2017;62(10):2915-22.
2. Fitzpatrick E, Dhawan A. Paediatric NAFLD: A distinct disease with the propensity for progressive fibrosis. . In: Williams R, Taylor-Robinson SD, editors. *Clinical Dilemmas in Non-Alcoholic Fatty Liver Disease* Wiley-Blackwell; 2016. p. 29-35.
3. Poston L, Caleyachetty R, Cnattingius S, Corvalan C, Uauy R, Herring S, et al. Preconceptional and maternal obesity: epidemiology and health consequences. *Lancet Diabetes Endocrinol*. 2016;4(12):1025-36.
4. Schwimmer JB, Deutsch R, Kahen T, Lavine JE, Stanley C, Behling C. Prevalence of fatty liver in children and adolescents. *Pediatrics*. 2006;118(4):1388-93.

5. Lawlor DA, Callaway M, Macdonald-Wallis C, Anderson E, Fraser A, Howe LD, et al. Nonalcoholic fatty liver disease, liver fibrosis, and cardiometabolic risk factors in adolescence: a cross-sectional study of 1874 general population adolescents. *J Clin Endocrinol Metab.* 2014;99(3):E410-7.
6. Xanthakos SA, Jenkins TM, Kleiner DE, Boyce TW, Mourya R, Karns R, et al. High Prevalence of Nonalcoholic Fatty Liver Disease in Adolescents Undergoing Bariatric Surgery. *Gastroenterology.* 2015;149(3):623-34 e8.
7. Anderson EL, Howe LD, Jones HE, Higgins JP, Lawlor DA, Fraser A. The Prevalence of Non-Alcoholic Fatty Liver Disease in Children and Adolescents: A Systematic Review and Meta-Analysis. *PLoS One.* 2015;10(10):e0140908.
8. Leung JC, Loong TC, Wei JL, Wong GL, Chan AW, Choi PC, et al. Histological severity and clinical outcomes of nonalcoholic fatty liver disease in nonobese patients. *Hepatology.* 2017;65(1):54-64.
9. Brumbaugh DE, Friedman JE. Developmental origins of nonalcoholic fatty liver disease. *Pediatr Res.* 2014;75(1-2):140-7.
10. Newton KP, Feldman HS, Chambers CD, Wilson L, Behling C, Clark JM, et al. Low and High Birth Weights Are Risk Factors for Nonalcoholic Fatty Liver Disease in Children. *J Pediatr.* 2017;187:141-6 e1.
11. Brumbaugh DE, Tearse P, Cree-Green M, Fenton LZ, Brown M, Scherzinger A, et al. Intrahepatic fat is increased in the neonatal offspring of obese women with gestational diabetes. *J Pediatr.* 2013;162(5):930-6 e1.
12. Modi N, Murgasova D, Ruager-Martin R, Thomas EL, Hyde MJ, Gale C, et al. The influence of maternal body mass index on infant adiposity and hepatic lipid content. *Pediatr Res.* 2011;70(3):287-91.
13. Patel KR, White FV, Deutsch GH. Hepatic steatosis is prevalent in stillborns delivered to women with diabetes mellitus. *J Pediatr Gastroenterol Nutr.* 2015;60(2):152-8.
14. Lager S, Gaccioli F, Ramirez VI, Jones HN, Jansson T, Powell TL. Oleic acid stimulates system A amino acid transport in primary human trophoblast cells mediated by toll-like receptor 4. *J Lipid Res.* 2013;54(3):725-33.
15. Deodati A, Argemi J, Germani D, Puglianiello A, Alisi A, De Stefanis C, et al. The exposure to uteroplacental insufficiency is associated with activation of unfolded protein response in postnatal life. *PLoS One.* 2018;13(6):e0198490.
16. Poston L. Maternal obesity, gestational weight gain and diet as determinants of offspring long term health. *Best Pract Res Clin Endocrinol Metab.* 2012;26(5):627-39.

17. Weng YH, Yang CY, Chiu YW. Risk Assessment of Adverse Birth Outcomes in Relation to Maternal Age. *PLoS One*. 2014;9(12):e114843.
18. Ayonrinde OT, Oddy WH, Adams LA, Mori TA, Beilin LJ, de Klerk N, et al. Infant nutrition and maternal obesity influence the risk of non-alcoholic fatty liver disease in adolescents. *J Hepatol*. 2017;67(3):568-76.
19. Kalliomaki M, Collado MC, Salminen S, Isolauri E. Early differences in fecal microbiota composition in children may predict overweight. *Am J Clin Nutr*. 2008;87(3):534-8.
20. Wankhade UD, Zhong Y, Kang P, Alfaro M, Chintapalli SV, Thakali KM, et al. Enhanced offspring predisposition to steatohepatitis with maternal high-fat diet is associated with epigenetic and microbiome alterations. *PLoS One*. 2017;12(4):e0175675.
21. Gemma C, Sookoian S, Alvarinas J, Garcia SI, Quintana L, Kanevsky D, et al. Maternal pregestational BMI is associated with methylation of the PPARGC1A promoter in newborns. *Obesity (Silver Spring)*. 2009;17(5):1032-9.
22. Aye IL, Rosario FJ, Powell TL, Jansson T. Adiponectin supplementation in pregnant mice prevents the adverse effects of maternal obesity on placental function and fetal growth. *Proc Natl Acad Sci U S A*. 2015;112(41):12858-63.
23. Lomas-Soria C, Reyes-Castro LA, Rodriguez-Gonzalez GL, Ibanez CA, Bautista CJ, Cox LA, et al. Maternal obesity has sex-dependent effects on insulin, glucose and lipid metabolism and the liver transcriptome in young adult rat offspring. *J Physiol*. 2018;596(19):4611-28.
24. McCurdy CE, Bishop JM, Williams SM, Grayson BE, Smith MS, Friedman JE, et al. Maternal high-fat diet triggers lipotoxicity in the fetal livers of nonhuman primates. *J Clin Invest*. 2009;119(2):323-35.
25. Romeo S, Kozlitina J, Xing C, Pertsemlidis A, Cox D, Pennacchio LA, et al. Genetic variation in PNPLA3 confers susceptibility to nonalcoholic fatty liver disease. *Nat Genet*. 2008;40(12):1461-5.
26. Romeo S, Sentinelli F, Cambuli VM, Incani M, Congiu T, Matta V, et al. The 148M allele of the PNPLA3 gene is associated with indices of liver damage early in life. *J Hepatol*. 2010;53(2):335-8.
27. Kozlitina J, Smagris E, Stender S, Nordestgaard BG, Zhou HH, Tybjaerg-Hansen A, et al. Exome-wide association study identifies a TM6SF2 variant that confers susceptibility to nonalcoholic fatty liver disease. *Nat Genet*. 2014;46(4):352-6.

28. Goffredo M, Caprio S, Feldstein AE, D'Adamo E, Shaw MM, Pierpont B, et al. Role of TM6SF2 rs58542926 in the pathogenesis of nonalcoholic pediatric fatty liver disease: A multiethnic study. *Hepatology*. 2016;63(1):117-25.
29. Chambers JC, Zhang W, Sehmi J, Li X, Wass MN, Van der Harst P, et al. Genome-wide association study identifies loci influencing concentrations of liver enzymes in plasma. *Nat Genet*. 2011;43(11):1131-8.
30. Wattacheril J, Lavine JE, Chalasani NP, Guo X, Kwon S, Schwimmer J, et al. Genome-Wide Associations Related to Hepatic Histology in Nonalcoholic Fatty Liver Disease in Hispanic Boys. *J Pediatr*. 2017;190:100-7 e2.
31. Jensen T, Abdelmalek MF, Sullivan S, Nadeau KJ, Green M, Roncal C, et al. Fructose and sugar: A major mediator of non-alcoholic fatty liver disease. *J Hepatol*. 2018;68(5):1063-75.
32. Mendez MA, Miles DR, Poti JM, Sotres-Alvarez D, Popkin BM. Persistent disparities over time in the distribution of sugar-sweetened beverage intake among children in the United States. *Am J Clin Nutr*. 2019;109(1):79-89.
33. Vandevijvere S, De Ridder K, Fiolet T, Bel S, Tafforeau J. Consumption of ultra-processed food products and diet quality among children, adolescents and adults in Belgium. *Eur J Nutr*. 2018.
34. Oddy WH, Herbison CE, Jacoby P, Ambrosini GL, O'Sullivan TA, Ayonrinde OT, et al. The Western dietary pattern is prospectively associated with nonalcoholic fatty liver disease in adolescence. *Am J Gastroenterol*. 2013;108(5):778-85.
35. Liu X, Peng Y, Chen S, Sun Q. An observational study on the association between major dietary patterns and non-alcoholic fatty liver disease in Chinese adolescents. *Medicine (Baltimore)*. 2018;97(17):e0576.
36. Gibson PS, Lang S, Gilbert M, Kamat D, Bansal S, Ford-Adams ME, et al. Assessment of Diet and Physical Activity in Paediatric Non-Alcoholic Fatty Liver Disease Patients: A United Kingdom Case Control Study. *Nutrients*. 2015;7(12):9721-33.
37. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014;384(9945):766-81.
38. Hegarty R, Deheragoda M, Fitzpatrick E, Dhawan A. Paediatric fatty liver disease (PeFLD): All is not NAFLD - Pathophysiological insights and approach to management. *J Hepatol*. 2018;68(6):1286-99.
39. Fitzpatrick E, Dhawan A. Noninvasive biomarkers in non-alcoholic fatty liver disease: current status and a glimpse of the future. *World J Gastroenterol*. 2014;20(31):10851-63.

40. Singh S, Venkatesh SK, Loomba R, Wang Z, Sirlin C, Chen J, et al. Magnetic resonance elastography for staging liver fibrosis in non-alcoholic fatty liver disease: a diagnostic accuracy systematic review and individual participant data pooled analysis. *Eur Radiol*. 2016;26(5):1431-40.
41. Hsu C, Caussy C, Imajo K, Chen J, Singh S, Kaulback K, et al. Magnetic Resonance vs Transient Elastography Analysis of Patients With Nonalcoholic Fatty Liver Disease: A Systematic Review and Pooled Analysis of Individual Participants. *Clin Gastroenterol Hepatol*. 2018.
42. Fitzpatrick E, Quaglia A, Vimalasvaran S, Basso MS, Dhawan A. Transient elastography is a useful noninvasive tool for the evaluation of fibrosis in paediatric chronic liver disease. *J Pediatr Gastroenterol Nutr*. 2013;56(1):72-6.
43. Eddowes PJ, Sasso M, Allison M, Tsochatzis E, Anstee QM, Sheridan D, et al. Accuracy of FibroScan Controlled Attenuation Parameter and Liver Stiffness Measurement in Assessing Steatosis and Fibrosis in Patients With Non-alcoholic Fatty Liver Disease. *Gastroenterology*. 2019.
44. Lee MS, Bae JM, Joo SK, Woo H, Lee DH, Jung YJ, et al. Prospective comparison among transient elastography, supersonic shear imaging, and ARFI imaging for predicting fibrosis in nonalcoholic fatty liver disease. *PLoS One*. 2017;12(11):e0188321.
45. Berna-Serna JD, Sanchez-Jimenez R, Velazquez-Marin F, Sainz de Baranda P, Guzman-Aroca F, Fernandez-Hernandez C, et al. Acoustic radiation force impulse imaging for detection of liver fibrosis in overweight and obese children. *Acta Radiol*. 2018;59(2):247-53.
46. Ishiba H, Sumida Y, Tanaka S, Yoneda M, Hyogo H, Ono M, et al. The novel cutoff points for the FIB4 index categorized by age increase the diagnostic accuracy in NAFLD: a multi-center study. *J Gastroenterol*. 2018;53(11):1216-24.
47. Bedogni G, Bellentani S, Miglioli L, Masutti F, Passalacqua M, Castiglione A, et al. The Fatty Liver Index: a simple and accurate predictor of hepatic steatosis in the general population. *BMC Gastroenterol*. 2006;6:33.
48. Alkhouri N, Mansoor S, Giammaria P, Liccardo D, Lopez R, Nobili V. The development of the pediatric NAFLD fibrosis score (PNFS) to predict the presence of advanced fibrosis in children with nonalcoholic fatty liver disease. *PLoS One*. 2014;9(8):e104558.
49. Vajro P, Lenta S, Socha P, Dhawan A, McKiernan P, Baumann U, et al. Diagnosis of nonalcoholic fatty liver disease in children and adolescents: position paper of the ESPGHAN Hepatology Committee. *J Pediatr Gastroenterol Nutr*. 2012;54(5):700-13.



50. Vos MB, Abrams SH, Barlow SE, Caprio S, Daniels SR, Kohli R, et al. NASPGHAN Clinical Practice Guideline for the Diagnosis and Treatment of Nonalcoholic Fatty Liver Disease in Children: Recommendations from the Expert Committee on NAFLD (ECON) and the North American Society of Pediatric Gastroenterology, Hepatology and Nutrition (NASPGHAN). *J Pediatr Gastroenterol Nutr.* 2017;64(2):319-34.
51. Molleston JP, Schwimmer JB, Yates KP, Murray KF, Cummings OW, Lavine JE, et al. Histological abnormalities in children with nonalcoholic fatty liver disease and normal or mildly elevated alanine aminotransferase levels. *J Pediatr.* 2014;164(4):707-13 e3.
52. Fitzpatrick E, Dew TK, Quaglia A, Sherwood RA, Mitry RR, Dhawan A. Analysis of adipokine concentrations in paediatric non-alcoholic fatty liver disease. *Pediatr Obes.* 2012;7(6):471-9.
53. Kleiner DE, Brunt EM, Van Natta M, Behling C, Contos MJ, Cummings OW, et al. Design and validation of a histological scoring system for nonalcoholic fatty liver disease. *Hepatology.* 2005;41(6):1313-21.
54. Angulo P, Kleiner DE, Dam-Larsen S, Adams LA, Bjornsson ES, Charatcharoenwitthaya P, et al. Liver Fibrosis, but No Other Histologic Features, Is Associated With Long-term Outcomes of Patients With Nonalcoholic Fatty Liver Disease. *Gastroenterology.* 2015;149(2):389-97 e10.
55. Brunt EM, Kleiner DE, Wilson LA, Unalp A, Behling CE, Lavine JE, et al. Portal chronic inflammation in nonalcoholic fatty liver disease (NAFLD): a histologic marker of advanced NAFLD- Clinicopathologic correlations from the nonalcoholic steatohepatitis clinical research network. *Hepatology.* 2009;49(3):809-20.
56. Gebhardt R, Matz-Soja M. Liver zonation: Novel aspects of its regulation and its impact on homeostasis. *World J Gastroenterol.* 2014;20(26):8491-504.
57. Nobili V, Mosca A, De Vito R, Raponi M, Scorletti E, Byrne CD. Liver zonation in children with non-alcoholic fatty liver disease: Associations with dietary fructose and uric acid concentrations. *Liver Int.* 2018;38(6):1102-9.
58. Feldstein AE, Charatcharoenwitthaya P, Treeprasertsuk S, Benson JT, Enders FB, Angulo P. The natural history of non-alcoholic fatty liver disease in children: a follow-up study for up to 20 years. *Gut.* 2009;58(11):1538-44.
59. Goyal NP, Schwimmer JB. The Progression and Natural History of Pediatric Nonalcoholic Fatty Liver Disease. *Clin Liver Dis.* 2016;20(2):325-38.

60. Xanthakos S, Lavine J, Yates K, Schwimmer JB, Molleston J, Rosenthal N, et al. Natural history of non alcoholic fatty liver disease in children receiving standard lifestyle counselling and placebo in NASH Clinical Research Network trials. *Hepatology*. 2017;66(1(supp)):31A.
61. Nobili V, Marcellini M, Devito R, Ciampalini P, Piemonte F, Comparcola D, et al. NAFLD in children: a prospective clinical-pathological study and effect of lifestyle advice. *Hepatology*. 2006;44(2):458-65.
62. DeVore S, Kohli R, Lake K, Nicholas L, Dietrich K, Balistreri WF, et al. A multidisciplinary clinical program is effective in stabilizing BMI and reducing transaminase levels in pediatric patients with NAFLD. *J Pediatr Gastroenterol Nutr*. 2013;57(1):119-23.
63. Musso G, Cassader M, Rosina F, Gambino R. Impact of current treatments on liver disease, glucose metabolism and cardiovascular risk in non-alcoholic fatty liver disease (NAFLD): a systematic review and meta-analysis of randomised trials. *Diabetologia*. 2012;55(4):885-904.
64. Vilar-Gomez E, Martinez-Perez Y, Calzadilla-Bertot L, Torres-Gonzalez A, Gra-Oramas B, Gonzalez-Fabian L, et al. Weight Loss Through Lifestyle Modification Significantly Reduces Features of Nonalcoholic Steatohepatitis. *Gastroenterology*. 2015;149(2):367-78 e5; quiz e14-5.
65. Lavine JE, Schwimmer JB, Van Natta ML, Molleston JP, Murray KF, Rosenthal P, et al. Effect of vitamin E or metformin for treatment of nonalcoholic fatty liver disease in children and adolescents: the TONIC randomized controlled trial. *JAMA*. 2011;305(16):1659-68.
66. Schwimmer JB, Lavine JE, Wilson LA, Neuschwander-Tetri BA, Xanthakos SA, Kohli R, et al. In Children With Nonalcoholic Fatty Liver Disease, Cysteamine Bitartrate Delayed Release Improves Liver Enzymes but Does Not Reduce Disease Activity Scores. *Gastroenterology*. 2016;151(6):1141-54 e9.
67. Singh T, Kochhar GS, Goh GB, Schauer P, Brethauer S, Kroh M, et al. Safety and efficacy of bariatric surgery in patients with advanced fibrosis. *Int J Obes (Lond)*. 2017;41(3):443-9.

**Table 1: Differential diagnosis of fatty liver in children and young people.**

Differential diagnosis	Diagnosis
Wilson disease	Low ceruloplasmin, high urinary or tissue copper, mutational analysis
Alpha 1 antitrypsin deficiency	Phenotype ?Genotype
Drugs – steroids, amiodarone, alcohol, methotrexate, MDMA (ecstasy), l-asparaginase, vitamin E, valproate, tamoxifen, antiretrovirals	History
Cystic Fibrosis associated liver disease	History / sweat test or mutational analysis
Malnutrition	History
Coeliac disease	Tissue transglutaminase / IgA, HLA typing, jejunal biopsy
Hepatitis C	HCV antibody status
Parenteral Nutrition associated liver disease	History
Mitochondrial disease / fatty acid oxidase deficiency	Lactate, acylcarnitines, respiratory chain enzymes, mutational analysis
Metabolic disease: Lysosomal acid lipase deficiency (Cholesterol Ester Storage Disease)	White cell enzymes, mutational analysis
Galactosaemia	Gal-1-PUT
Fructosaemia	Enzymology
Glycogen storage disease	White cell enzymes, mutational analysis
Peroxisomal disorders	Very long chain fatty acids, mutational analysis
Mauriac syndrome	History of type 1 diabetes
Hypobetalipoproteinaemia/ abetalipoproteinaemia	Low lipid levels, reduced / absent Apo1B, \ mutational analysis
Lipodystrophies	Mutational analysis
Schwachman syndrome	Pancreatic insufficiency / mutational analysis